

TOE AND METATARSAL FRACTURES



American College of
Foot and Ankle Surgeons

The structure of your foot is complex, consisting of bones, muscles, tendons, and other soft tissues. Of the 26 bones in your foot, 19 are toe bones (phalanges) and metatarsal bones (the long bones in the midfoot). Fractures of the toe and metatarsal bones are common and require evaluation by a specialist. A podiatric foot and ankle surgeon should be seen for proper diagnosis and treatment, even if initial treatment has been received in an emergency room.


What Is a Fracture?

A fracture is a break in the bone. Fractures can be divided into two categories: traumatic fractures and stress fractures.

Traumatic fractures (also called acute fractures) are caused by a direct blow or impact—like seriously stubbing your toe. Traumatic fractures can be *displaced* or *nondisplaced*. If the fracture is displaced, the bone is broken in such a way that it has changed in position (dislocated). Treatment of a traumatic fracture depends on the location and extent of the break and whether it is displaced. Surgery is sometimes required.

Signs and symptoms of a traumatic fracture include:

- You may hear a sound at the time of the break.
- “Pinpoint pain” (pain at the place of impact) at the time the fracture occurs and perhaps for a few hours later, but often the pain goes away after several hours.



- Deviation (misshapen or abnormal appearance) of the toe.
- Bruising and swelling the next day.
- It is not true that “if you can walk on it, it’s not broken.” Evaluation by the podiatric surgeon is always recommended.

Stress fractures are tiny, hairline breaks that are usually caused by repetitive stress. Stress fractures often afflict athletes who, for example, too rapidly increase their running mileage. Or they may be caused by an abnormal foot structure, deformities, or osteoporosis. Improper footwear may also lead to stress fractures.

Stress fractures should not be ignored, because they will come back unless properly treated. Symptoms of stress fractures include:

- Pain with or after normal activity

- Pain that goes away when resting and then returns when standing or during activity
- “Pinpoint pain” (pain at the site of the fracture) when touched
- Swelling, but no bruising

Sprains and fractures have similar symptoms, although sometimes with a sprain, the whole area hurts rather than just one point. Your podiatric surgeon will be able to diagnose which you have and provide appropriate treatment. Certain sprains or dislocations can be severely disabling. Without proper treatment they can lead to crippling arthritis.

Consequences of Improper Treatment

Some people say that “the doctor can’t do anything for a broken bone in the foot.” This is usually not true. In fact, if a fractured toe or metatarsal bone is not treated correctly, serious complications may develop. For example:

- A deformity in the bony architecture which may limit the ability to move the foot or cause difficulty in fitting shoes
- Arthritis, which may be caused by a fracture in a joint (the juncture where two bones meet), or may be a result of angular deformities that develop when a displaced fracture is severe or hasn’t been properly corrected
- Chronic pain and long-term dysfunction

- Non-union, or failure to heal, can lead to subsequent surgery or chronic pain.

Treatment of Toe Fractures

Fractures of the toe bones are almost always traumatic fractures. Treatment for traumatic fractures depends on the break itself and may include these options:

- **Rest.** Sometimes rest is all that is needed to treat a traumatic fracture of the toe.
- **Splinting.** The toe may be fitted with a splint to keep it in a fixed position.
- **Rigid or stiff-soled shoe.** Wearing a stiff-soled shoe protects the toe and helps keep it properly positioned.
- **"Buddy taping"** the fractured toe to another toe is sometimes appropriate, but in other cases it may be harmful.
- **Surgery.** If the break is badly displaced or if the joint is affected, surgery may be necessary. Surgery often involves the use of fixation devices, such as pins.

Treatment of Metatarsal Fractures


Breaks in the metatarsal bones may be either stress or traumatic fractures. Certain kinds of fractures of the metatarsal bones present unique challenges.

For example, sometimes a fracture of the first metatarsal bone (behind the big toe) can lead to arthritis. Since the big toe is used so frequently and bears more weight than other toes, arthritis in that area can make it painful to walk, bend, or even stand.

Another type of break, called a Jones fracture, occurs at the base of the fifth metatarsal bone (behind the little toe). It is often misdiagnosed as an ankle sprain, and misdiagnosis can have serious consequences since sprains and fractures require different treatments. Your podiatric surgeon is an expert in correctly identifying these conditions as well as other problems of the foot.

Treatment of metatarsal fractures depends on the type and extent of the fracture, and may include:

- **Rest.** Sometimes rest is the only treatment needed to promote healing of a stress or traumatic fracture of a metatarsal bone.
- **Avoid the offending activity.** Because stress fractures result from repetitive stress, it is important to avoid the activity that led to the fracture. Crutches or a wheelchair are sometimes required to offload weight from the foot to give it time to heal.
- **Immobilization, casting, or rigid shoe.** A stiff-soled shoe or other form of immobilization may be used to protect the fractured bone while it is healing.
- **Surgery.** Some traumatic fractures of the metatarsal bones require surgery, especially if the break is badly displaced.
- **Follow-up care.** Your podiatric foot and ankle surgeon will provide instructions for care following surgical or non-surgical treatment. Physical therapy, exercises and rehabilitation may be included in a schedule for return to normal activities. ▲



**American College of
Foot and Ankle Surgeons**

This information has been prepared by the Consumer Education Committee of the American College of Foot and Ankle Surgeons, a professional society of 5,700 podiatric foot and ankle surgeons. Members of the College are Doctors of Podiatric Medicine who have received additional training through surgical residency programs.

The mission of the College is to promote superior care of foot and ankle surgical patients through education, research and the promotion of the highest professional standards.

Copyright © 2004, American College of Foot and Ankle Surgeons • www.acfas.org